INDEX

Acorns, feeding of, to laying pullets and ducks, 175–81; experimental, 177–8; tannin-content of, and changes on boiling, 178–9; yolk-discoloration of eggs, 175 sqq.

Banana, genetic system of varieties of, in relation to banana breeding, 89–98; breeding programme, 94–8; parthenocarpy, 90–1, 93–4; polyploidy, 92–3; sterility, 91–2; vegetative propagation, 89–90.
Barlow, H. W. B., *see* Crowther, F.
Beef cattle in peace and war, 191–228; composition of body, 196–8; cost of growing and fattening, 207–21; dietetic qualities of beef, 194–6; effect of food on development of a beef steer, 227–8; efficiency of conversion into human food, 226; experimental data, Minnesota, 192 sqq., Missouri, 193 sqq.; optimum slaughtering point, 221–4; valuation of beef constituents, 224–6; yield of human food, 198–207.
Brett, C. E., *see* Lovatt, J.

Clove trees, stem-girth as criterion of canopy-area and yield, 33–7.
Coffee, control of a mealy bug on, in Kenya, 78–88.
Common, R. H., the feeding-value of flax by-products, 229–33.
Cotton, efficiency of progeny-row-breeding in improving, 140–54; design of a breeding project, 152–3; genetic basis of progeny selection, 149–52; Variety U4, effect of selection on, 147–9; Variety V135, effect of selection on, 141–7, history of, 140–1, 149.
— influence of weeds on, in Sudan Gezira, 1–14; clean weeding of the crop, 2–6; effect of weeding land resting before cotton, 6–10; effect of nitrogen supply, 11–13; effect of termites, 11.
— tap-root damage ascribed to termites, 99–112.
Cowie, G. A., the relative responses of the potato crop to different potash fertilizers, 23–32.
Crowther, F., influence of weeds on cotton in the Sudan Gezira, 1–14.
— and H. W. B. Barlow, tap-root damage of cotton, ascribed to termites, in the Sudan Gezira, 99–112.

Deli Tobacco Experiment Station (Sumatra), research at, 15–22; agrochemical research, 21–2; insect pests, 18–21; manuring, 15–16, 21–2; plant diseases, 17–18; slime disease, 16–17.
Diseases and pests: of coffee in Kenya, 78–88; potatoes, 126 sqq.; tobacco in Sumatra, 16, 17–21.
Dodds, K. S., the genetic system of banana varieties in relation to banana breeding, 89–98.
Ducks, laying, feeding of acorns to, 175–81.

Fertilizers and manures: for mangolds, 49–77; potatoes, 23–32; tobacco, 15–16.
Flax by-products, feeding-value of, 229–33; flax chaff, 229, 230, 231–2, and fibre, 232; linseed and 'feeding linseed', 229, 230; shives, &c., 230, 232; whole flax bolls, 229, 230.

Hutchinson, J. B., the efficiency of progeny-row-breeding in cotton improvement, 140–54.

Kenya, mealy bug on coffee and other crops in, 78–88.

Le Pelley, R. H., the biological control of a mealy bug on coffee and other crops in Kenya, 78–88.
Lovatt, J., fattening of pigs on swill alone, 182–90.

Mangolds, the Rothamsted experiments on, 1872–1940, 49–77. Pt. I. Effect of manures on yield of roots, 49–64: Barnfield expt., 50–3; effects of different forms of nitrogen, 62–4, of inorganic fertilizers, 53–7, of organic manures, 58–62; historical, 49–50. Pt. II. Effect of manures on plant-growth, 65–77; bolting, 74, 75–6; effects on competition, plant-number, survival, yield, 65–70; on the soil, 76–7; spacing, 70–2; yield of leaves, 72–4.
Manning, H. L., *see* Hutchinson, J. B.
Mealy bug, biological control of a, on coffee and other crops in Kenya, 78–88; first attempts, 78–9; later attempts, 80–1; liberations since 1938, 82–3; parasites for, from Uganda, 79–80; position before introduction from Uganda, 80–2, after establishment, 83–8.
Milton, W. E. J., the buried viable-seed content of a Midland calcareous clay soil, 155–66 [and two insets].

Pasture, value of, for meal-fed pigs, 168–74; experimental method, 169–70; results, 170–4.
Pasture herbage, effects of defoliation intensity and fertilizer treatment on Transvaal Highveld, 113–24; experimental, 114–15; herbage, composition of, 117–18, yields, 115–17; roots, composition of, 119–21; storage and exhaustion of reserves, 121–2, weights, 118–19.
Pickup, J., *see* Lovatt, J.
Pigs, fattening of, on swill alone, 182–90; carcass and bacon, 185–7; disease, 187–8; financial aspect, 187; market for bacon and pork, 189; the stock, 183, 185, 187–8; the swill, 182–3, 184, 188–9.
Plant breeding: banana, 89–98; cotton, 140–54; potato, 125–39; sugar-cane, 38–48.

Potato breeding, recent research in, 125–39; blight-resistant varieties, 131, 132; future progress, 133; mutability of the tuber, 129–30; pedigree of new resistant variety, 131; recent Russian discoveries, 129, 133; varieties, transmissible characters of, and their sources, 133–8; virus disease, 126–8; virus-resistant varieties, 128–9; wart disease, 125–6.
Potatoes, response of, to potash fertilizers, 23–32; chlorine, 25–7; complete fertilizers, 24–5; composition, 27–9; cooking quality, 30–1; rainfall, 29–30.
Pullets, laying, feeding of acorns to, 175–81.

Roelofsen, P. A., recent research at the Deli Tobacco Research Station, Medan, Sumatra, 15–22.
Russell, Sir E. J., *see* Watson, D. J.

Salaman, R. N., recent research in potato breeding, 125–39.
Seed, buried, viable, in a Midland calcareous soil, 155–67; experimental method, 156; germination, 161, 162; grass and arable fields at Dodwell and Drayton, 157 sqq.; leguminous species, 162; soil sampling, 156, 162–3.
Stevenson, G. C., the present and potential value of sugar-cane breeding, 38–48.
Sudan (Anglo-Egyptian), influence of weeds on cotton in, 1–14; termite damage to cotton roots, 99–112.
Sugar-cane breeding, present and potential value of, 38–48; history, 39–40; problems, 40–4; trend of research, 44–8.

Temperton, H., the feeding of acorns to laying pullets and ducks, 175–81.
Termites, effect of, on tropical soils, 12.
— tap-root damage of cotton, ascribed to, 99–112; damage in relation to season and crop-rotation, 101–3, crop-yield, 107, cultural treatments, 105–7, manuring, 102–4, variety of cotton, 104.
Tidbury, G. E., stem-girth as a criterion in field trials with young clove trees, 33–7.
Tobacco, research in Sumatra, 15–22; diseases, 16–18; insect pests, 18–21; manuring, 15–16, 21–2; soils, 21.

Wallace, L. R., the value of pasture for meal-fed pigs, 168–74.
Watson, D. J., the Rothamsted experiments on mangolds, 1872–1940. Pt. I, 49–64. Pt. II, 65–77.
Watson, D. M. S., beef cattle in peace and war, 191–228.
Weeds, buried viable seeds of, in a calcareous soil, 155 sqq.
— effect of on cotton, 1–14; on prevention of soil erosion, 1, 11.
Weinmann, H., effects of defoliation intensity and fertilizer treatment on Transvaal Highveld, 113–24.
Worden, A. N., *see* Lovatt, J.

CORRIGENDA

Vol. X, No. 40, p. 221, par. 3, last 2 lines, *read* '. . . since the treatments were stopped at the end of 1935'.

Vol. XI, No. 42, p. 80, line 11, *read* '. . . the most effective one. Although, more theoretically . . .'

—— p. 81, line 36, *read* 'The crop loss over the mealy-bug area during this period is not thought to have been less than 5 per cent.'

—— p. 82, line 13, *read* '. . . many useful indigenous herbs, and shrubs . . .'

—— p. 85, column headed '1938' line 17, for '375' *read* '275'.

—— p. 100, lines 16 and 22, and Plate 1, legend, for (above) *read* (a), for (below) *read* (b).

—— p. 110, line 13, add as footnote: Unpublished note 1935–6: Larger form Termes (Cyclotermes) sp.; smaller form Microtermes thorocalis Sjöst. 'Mortality amongst young cotton plants was invariably caused by the latter, whereas the (wooden) pegs were attacked by both.'

PRINTED IN GREAT BRITAIN AT THE UNIVERSITY PRESS, OXFORD
BY JOHN JOHNSON, PRINTER TO THE UNIVERSITY